arXiv:1211.1083v2 [hep-ph] 8 Nov 2012

125 GeV Higgs from a not so light Technicolor Scalar

Roshan Foadi[♣], Mads T. Frandsen[♥], and Francesco Sannino[♥]

[♣] *Centre for Cosmology, Particle Physics and Phenomenology (CP3) Chemin du Cyclotron 2*
Université catholique de Louvain, Belgium and
[♥] CP^3-Origins *& the Danish Institute for Advanced Study* DIAS,
University of Southern Denmark, Campusvej 55, DK-5230 Odense M, Denmark.

Assuming that the observed Higgs-like resonance at the Large Hadron Collider is a technicolor isosinglet scalar (the technicolor Higgs), we argue that the standard model top-induced radiative corrections reduce its dynamical mass towards the desired experimental value.

We then discuss conditions for the spectrum of technicolor theories to feature a technicolor Higgs with the phenomenologically required dynamical mass. We use scaling laws coming from modifying the technicolor matter representation, number of technicolors, techniflavors as well as the number of doublets gauged under the electroweak theory. Finally we briefly summarize the potential effects of walking dynamics on the technicolor Higgs.

Preprint: CP^3-Origins-2012-028 & DIAS-2012-29

I. INTRODUCTION

Recently the ATLAS and CMS collaborations have announced the discovery of a new boson with the approximate mass of 125 GeV [1, 2]. The observed decays to standard model (SM) diboson pairs, $\gamma\gamma$, ZZ^* and WW^*, are in rough agreement with those expected from the SM Higgs. Here we will assume this state is a scalar as suggested by the observed decays into ZZ^* and WW^* [3–5].

If new strong dynamics similar to technicolor (TC) [6–8] underlies the Higgs mechanism a pressing question is: What makes this scalar light compared to the natural scale of the electroweak theory $\Lambda \sim 4\pi v$ with $v \simeq 246$ GeV ?

To answer this question we must disentangle the SM radiative corrections from the dynamical mass M_H^0, i.e. the mass stemming purely from composite dynamics. We will argue that the dynamical mass of the scalar required to match the observations is of the order of 600 GeV, which is substantially heavier than the observed value. This estimate assumes that Λ is the natural cutoff for the radiative corrections and that the scalar coupling to the top quark is similar to that of the SM Higgs. This is a significant increase in the value of the dynamical mass compared to the observed 125 GeV, often naively identified with the dynamical mass. In minimal TC models [9, 10] the compositeness scale is Λ.

Additionally, if the dominant decay channels are into SM states, then for a physical mass of 125 GeV the TC Higgs will be narrow simply because of kinematics. A similar example in strong dynamics is the $f_0(980)$, which is extremely narrow because the decay mode into $K\bar{K}$ is below threshold [11].

We first analyze the SM radiative corrections to the TC Higgs mass. We then discuss the scaling of the TC Higgs dynamical mass as a function of the number of techniflavors $N_{\rm TF}$, technicolors $N_{\rm TC}$, as well as the representation $R_{\rm TC}$ with respect to the TC gauge group. As reference theory we use 2-flavor QCD leading, which leads to a dynamical mass in the 1.0 TeV $\lesssim M_H^0 \lesssim$ 1.4 TeV range, before changing the *geometry* of the TC gauge theory. For example, we shall see that a TC Higgs becomes lighter as $d(R_{\rm TC})$ and/or $N_{\rm TF}^*$ grow.

A second source of reduction of the TC Higgs dynamical mass may come from walking dynamics [12]. The latter is useful to alleviate the tension with respect to the flavor changing neutral current problem and to reduce the value of the S-parameter [13, 14]. However, the S-parameter is not expected to vanish in conformal field theories [15–17] and the most recent study of precision data including a TC Higgs at the effective Lagrangian level appeared in [18]. In the literature such a light TC Higgs is also known as techni-dilaton [19–22].

The paper is organized as follows: In Sec. II we set up an effective Lagrangian including the TC Higgs, the SM particles, and their interactions. We then compute the corrections to the squared TC Higgs mass. In Sec. III we analyze the scaling of the dynamical TC Higgs mass as a function of $d(R_{\rm TC})$ and $N_{\rm TF}$, and show – taking into account the SM radiative corrections – which TC theories can give a physical TC Higgs mass of 125 GeV. We then turn to the possibility of further reductions of the dynamical mass from walking dynamics. We argue that walking TC can accommodate a 125 GeV Higgs even for small values of $d(R_{\rm TC})$ and number of technidoublets. Finally in Sec. IV we offer our conclusions.

II. RADIATIVE CORRECTIONS TO THE MASS OF THE TC HIGGS

We consider TC theories featuring at scales below the mass of the technirho M_ρ only the eaten Goldstone bosons and the TC Higgs (H). We assume the TC dynamics to respect the $SU(2)_c$ custodial isospin symmetry, and adopt a nonlinear realization for the composite states. The latter are thus classified according to linear multiplets of $SU(2)_c$: the electroweak Goldstone bosons π^a, with $a = 1, 2, 3$, form an $SU(2)_c$ triplet, whereas the TC Higgs is an $SU(2)_c$ singlet. The elementary SM fields are as linear multiplets of the electroweak group. The Yukawa interactions of the TC Higgs with SM fermions are induced by interactions beyond the TC theory itself, *e.g.* by extended TC (ETC) interactions [7, 8].

Assuming that the only non-negligible sources of custodial isospin violation are due to the Yukawa interactions, and retaining only the leading order terms in a momentum expansions, leads to the effective Lagrangian

$$\begin{aligned}\mathcal{L} = {} & \mathcal{L}_{\overline{\rm SM}} + \left(1 + \frac{2r_\pi}{v}H + \frac{s_\pi}{v^2}H^2\right)\frac{v^2}{4}{\rm Tr}\, D_\mu U^\dagger D^\mu U + \frac{1}{2}\,\partial_\mu H\,\partial^\mu H - V[H] \\ & - \; m_t\left(1 + \frac{r_t}{v}H\right)\left[\overline{q}_L\, U\left(\frac{1}{2} + T^3\right)q_R + {\rm h.c.}\right] \\ & - m_b\left(1 + \frac{r_b}{v}H\right)\left[\overline{q}_L\, U\left(\frac{1}{2} - T^3\right)q_R + {\rm h.c.}\right] + \cdots \\ & - \Delta S\; W^a_{\mu\nu}B^{\mu\nu}\;{\rm Tr}\; T^a U T^3 U^\dagger + O\left(\frac{1}{M_\rho}\right)\end{aligned} \qquad (1)$$

where $\mathcal{L}_{\overline{\rm SM}}$ is the SM Lagrangian without Higgs and Yukawa sectors, the ellipses denote Yukawa interactions for SM fermions other than the top-bottom doublet $q \equiv (t, b)$, and $O(1/M_\rho)$ includes higher-dimensional operators, which are suppressed by powers of $1/M_\rho$. In this Lagrangian $v \simeq$ 246 GeV is the electroweak vev, U is the usual exponential map of the Goldstone bosons produced



FIG. 1: Quadratically divergent diagrams contributing to the Higgs mass, with the interaction vertices given by (2). The gauge boson exchanges are computed in Landau gauge: then the seagull diagrams, with a single W and Z exchange, are the only quadratically divergent one-loop diagrams with gauge boson exchanges.

by the breaking of the electroweak symmetry, $U = \exp\left(i\pi^a T^a/v\right)$, with covariant derivative $D_\mu U \equiv \partial_\mu U - igW^a_\mu T^a U + ig' U B_\mu T^3$, $2T^a$ are the Pauli matrices, with $a = 1, 2, 3$, and $V[H]$ is the TC Higgs potential. ΔS is the contribution to the S parameter from the physics at the cutoff scale, and is assumed to vanish in the $M_\rho \to \infty$ limit. The interactions contributing to the Higgs self-energy are

$$\begin{aligned}\mathcal{L}_H \supset & \frac{2\, m_W^2\, r_\pi}{v}\, H\, W^+_\mu\, W^{-\mu} + \frac{m_Z^2\, r_\pi}{v}\, H\, Z_\mu\, Z^\mu - \frac{m_t\, r_t}{v}\, H\, \bar{t}\, t \\ & + \frac{m_W^2\, s_\pi}{v^2}\, H^2\, W^+_\mu\, W^{-\mu} + \frac{m_Z^2\, s_\pi}{2\, v^2}\, H^2\, Z_\mu\, Z^\mu\, . \end{aligned} \tag{2}$$

The tree-level SM is recovered for

$$r_\pi = s_\pi = r_t = r_b = 1\, . \tag{3}$$

We divide the radiative corrections to the TC Higgs mass into two classes: external contributions, corresponding to loop corrections involving elementary SM fields, and TC contributions, corresponding to loop corrections involving TC composites only. The latter contribute to the dynamical mass M_H^0, whose size will be estimated in the next section by non-perturbative analysis. In order to isolate the SM contributions we work in Landau gauge. Here transversely polarized gauge boson propagators correspond to elementary fields, and massless Goldstone boson propagators correspond to TC composites. The only SM contributions to the TC Higgs mass which are quadratically divergent in the cutoff come from the diagrams of Fig. 1. Retaining only the quadratically divergent terms leads to a physical mass M_H given by

$$M_H^2 = (M_H^0)^2 + \frac{3(4\pi\kappa F_\Pi)^2}{16\pi^2 v^2}\left[-4r_t^2 m_t^2 + 2s_\pi\left(m_W^2 + \frac{m_Z^2}{2}\right)\right] + \Delta_{M_H^2}(4\pi\kappa F_\Pi)\, , \tag{4}$$

where $\Delta_{M_H^2}(4\pi\kappa F_\Pi)$ is the scale-dependent counterterm and κ is a order unity number. To be able to provide a physical estimate we assume that the counterterm is negligible at the scale $4\pi\kappa F_\Pi$, where F_Π is the TC pion decay constant and κ scales like $1/\sqrt{d(R_{\rm TC})}$ if the cutoff is identified with the technirho mass, or is a constant if the cutoff is of the order of $4\pi F_\Pi$. Provided r_t is also of order one, the dominant radiative correction is due to the top quark. For instance, if $F_\Pi = v$, which is appropriate for a TC theory with one weak technidoublet, then $\delta M_H^2 \sim -12\kappa^2 r_t^2 m_t^2 \sim -\kappa^2 r_t^2 (600\,{\rm GeV})^2$. This demonstrates that the dynamical mass of the TC Higgs can be substantially heavier than the physical mass, $M_H \simeq 125$ GeV.

III. THE DYNAMICAL MASS OF THE TC HIGGS

In QCD the lightest scalar is the σ meson (also termed $f_0(500)$ in PDG), with a measured mass between 400 and 550 MeV [23] in agreement with early determinations [11]. Scaling up two-flavor QCD yields a TC Higgs dynamical mass in the 1.0 TeV $\lesssim M_H^0 \lesssim$ 1.4 TeV range. This estimate changes when considering TC theories which are not an exact replica of two-flavor QCD. Here we determine the *geometric scaling* of the TC Higgs dynamical mass, i.e. the value of M_H^0 as function of the TC matter representation $d(R_{\rm TC})$, $N_{\rm TC}$ and the number of techniflavors $N_{\rm TF}$ for a given $SU(N_{\rm TC})$ gauge theory. For a generalization to different gauge groups see [24, 25]. We then discuss possible effects of walking dynamics on M_H^0, which are not automatically included in the geometric scaling. Taking into account the SM induced radiative corrections discussed in Sec. II, we argue that TC can accommodate a TC Higgs with a physical mass of 125 GeV, with or without effects from walking.

A. Geometric Scaling of the TC Higgs mass

We will consider at most two-index representations for TC matter, since at large $N_{\rm TC}$ even higher representations loose quickly asymptotic freedom [26]. The relevant scaling rules are:

$$F_\Pi^2 \sim d(R_{\rm TC})\, m_{\rm TC}^2\,, \quad v^2 = N_{\rm TD}\, F_\Pi^2\,, \tag{5}$$

where F_Π is the technipion decay constant, $m_{\rm TC}$ is the dynamically generated constituent techniquark mass, and $N_{\rm TD} = N_{\rm TF}^*/2$, where $N_{\rm TF}^*$ is the actual number of techniflavors arranged in weak doublets and therefore $N_{\rm TF}^* \leq N_{\rm TF}$. $v = 246$ GeV is the electroweak vacuum expectation value and will be kept fix in the following.

The squared mass of any large $N_{\rm TC}$ leading technimeson scales like:

$$(M_H^0)^2 = \frac{3}{d(R_{\rm TC})}\,\frac{1}{N_{\rm TD}}\,\frac{v^2}{f_\pi^2}\, m_\sigma^2\,. \tag{6}$$

$SU(N_{\rm TC})$ Rep.	$d(R_{\rm TC})$	$N_{\rm TD}\ (M_H^0)^2$
$\square$	$N_{\rm TC}$	$\frac{v^2}{f_\pi^2}\left[\frac{3}{N_{\rm TC}}\right]^{1-p} m_\sigma^2$
$\begin{array}{c}\square\\[-4pt]\square\end{array}$	$\frac{N_{\rm TC}(N_{\rm TC}-1)}{2}$	$\frac{v^2}{f_\pi^2}\,\frac{3}{N_{\rm TC}(N_{\rm TC}-1)/2}\, m_\sigma^2$
$\square\!\square$	$\frac{N_{\rm TC}(N_{\rm TC}+1)}{2}$	$\frac{v^2}{f_\pi^2}\,\frac{3}{N_{\rm TC}(N_{\rm TC}+1)/2}\, m_\sigma^2$

TABLE I: Scaling formulae for M_H^0 obtained by scaling up the mass of the QCD σ meson. The general formula is given in (6), and here is applied to the F representation, with $p = 0$, the A representation, and the S representation.

The leading states for the fundamental (F) representation are fermion-antifermion pairs and in the case of the two-index representations are mesons containing any number of fermions [9, 27]. The normalization to three colors QCD can also be assumed for the two-index representations since the two-index antisymmetric (A) for three colors is exactly QCD and the two-index symmetric (S) at infinite number of TC colors cannot be distinguished from the antisymmetric one.

The mesonic states which are not leading at large $N_{\rm TC}$ for the fundamental representation will decouple from the leading ones. Their scaling is

$$(M_H^0)^2 = \left[\frac{N_{\rm TC}}{3}\right]^p \frac{3}{N_{\rm TC}}\,\frac{1}{N_{\rm TD}}\,\frac{v^2}{f_\pi^2}\, m_\sigma^2\,, \tag{7}$$

with $p > 0$ [9, 27]. Applying these general results to the F and the A representation gives the M_H^0 scaling formulae summarized in Tab. I.

The geometric scaling above can be compared to the dynamical mass of the TC Higgs required to fit the experiments once the electroweak corrections have been subtracted. Using (4) gives

$$N_{\rm TD}\ (M_H^0)^2 = N_{\rm TD}\ M_H^2 + 12\ \kappa^2 r_t^2 m_t^2 - 6\ \kappa^2 s_\pi \left(m_W^2 + \frac{m_Z^2}{2}\right)\,. \tag{8}$$

The top contribution dominates for $\mathcal{O}(1)$ values of $N_{\rm TD}$ and κr_t. With this assumption the contribution of the gauge bosons can be neglected. Using (6) we estimate κr_t, up to corrections due to the weak gauge boson exchange, to be

$$\kappa^2 r_t^2 = \frac{A}{B}\frac{1}{d(R_{\rm TC})} - \frac{N_{\rm TD}}{B} \qquad \text{with} \qquad A = 3\frac{v^2}{M_H^2}\frac{m_\sigma^2}{f_\pi^2}\,, \qquad B = 12\frac{m_t^2}{M_H^2}\,. \tag{9}$$

For the case in which κ is taken to scale like $\hat{\kappa}/\sqrt{d(R_{\rm TC})}$ we get

$$\hat{\kappa}^2 r_t^2 = \frac{A}{B} - \frac{N_{\rm TD}\,d(R_{\rm TC})}{B}\,. \tag{10}$$

For example, choosing $N_{\rm TD} = 1$ and the S representation with $N_{TC} = 3$ we have that $d(R_{\rm TC}) = 6$ and therefore using (9,10) we get $\kappa r_t \simeq 1.5$ ($\hat{\kappa} r_t \simeq 3.8$).

In figure 2 we plot the required dynamical mass $\sqrt{N_{\rm TD}}M_H^0$ to fit the observed resonance value for different values of κr_t. Since the dependence on $N_{\rm TD}$ is small we will show only the case of $N_{\rm TD} = 1$ and neglect, by setting $s_\pi = 0$, the weak gauge boson corrections. We choose $\kappa r_t = 0$ (dashed curve), $\kappa r_t = 1$ (solid curve), and $\kappa r_t = 1.5$ (dotted curve). The estimates of Tab. I are shown as the colored bands in the same figure, with the lower (upper) edges of the bands corresponding to the experimental lower (upper) bound on the QCD m_σ. The horizontal band is for the F representation and $p = 1$, the blue (middle) band for the A representation, and the red (lower) band for the S representation. From Fig. 2 we observe that if we *do* take the external radiative corrections into account, and set $\kappa r_t = 1$ ($\kappa r_t = 1.5$), then the geometric TC scaling can accommodate a 125 GeV Higgs. This occurs where the bands overlap with the middle solid (upper dotted) horizontal line. This requires $5 \le N_{\rm TC} \le 6$ ($2 \le N_{\rm TC} \le 4$) for the S representation, and $7 \le N_{\rm TC} \le 10$ ($4 \le N_{\rm TC} \le 6$) for the A representation. On the other hand, If we do not take the external radiative corrections into account, a TC theory with geometric scaling can only accommodate a 125 GeV Higgs where the bands overlap with the lower dashed horizontal line, requiring large values of $N_{\rm TC} \gtrsim 20$ for the S and A representation.

We conclude that radiative SM corrections cannot be neglected when discussing extensions of the SM featuring TC Higgs states, or any other extension similar to TC. Furthermore these corrections tend to reduce the physical mass allowing for more natural values of the underlying dynamical mass of the TC isosinglet scalar.

B. Walking effects on the TC Higgs mass

In models with walking dynamics the above scalings are expected to overestimate the mass of the TC Higgs. In fact, as $N_{\rm TC}$ and/or $N_{\rm TD}$ take on different values, gauge dynamics is expected to change. For a given TC gauge group and a given representation for the technifermions, there



FIG. 2: Estimates of the dynamical TC Higgs mass M_H^0. The bands are constructed using the geometric scaling of Tab. I, with gray, blue, and red band for F, A, and S representation, respectively. In each band the lower (upper) curve corresponds to the experimental lower (upper) bound on m_σ. The dashed, solid, and dotted curves show the required dynamical TC Higgs mass to achieve a 125 GeV Higgs, using (8) for $N_{\rm TD} = 1$, $s_\pi = 0$, and $\kappa r_t = 0$, $\kappa r_t = 1$, and $\kappa r_t = 1.5$, respectively.

exists a critical number of Dirac techniflavors, $N_{\rm TF}^c$, such that for $N_{\rm TF} < N_{\rm TF}^c$ chiral symmetry is broken, whereas for $N_{\rm TF} > N_{\rm TF}^c$ chiral symmetry is restored. For $N_{\rm TF}$ slightly above $N_{\rm TF}^c$, the gauge theory develops an infrared fixed point. Gauge dynamics becomes conformal at low energies, and the coupling never reaches the critical value for chiral symmetry breaking. For $N_{\rm TF}$ slightly below $N_{\rm TF}^c$, the TC force does break chiral symmetry, but may feel the presence of the nearby fixed point. In this case the theory becomes walking in a range of energies above the chiral symmetry breaking scale. Assuming a continuous phase transition, as $N_{\rm TF}$ approaches $N_{\rm TF}^c$ from below, the technihadron masses and decay constants approach zero [1]. It is reasonable to assume that for walking theories when $N_{\rm TF}$ is very close to $N_{\rm TF}^c$ one has [21]

$$(M_H^0)^2 \propto (N_{\rm TF}^c - N_{\rm TF})^{\nu_H} \frac{F_\Pi^2}{d(R_{\rm TC})} , \qquad (11)$$

---

[1] There is still the possibility that the phase transition is of *jumping* type [28, 29], meaning the transition is not continuous and therefore no massive state becomes light when approaching the conformal boundary. Jumping conformal phase transitions have been identified in [30]. It is relevant to mention that the first four-dimensional continuous conformal phase transition á la Miransky [31–33] as function of the number of flavors of the theory has been discovered in [34]. For interesting lower dimensional models displaying Miransky scaling investigated on the lattice we refer to [35, 36].

where ν_H depends on the specific underlying theory. Using the second of (5) to relate F_Π to v one deduces

$$N_{\rm TD}\,(M_H^0)^2 \propto (N_{\rm TF}^c - N_{\rm TF})^{\nu_H}\,\frac{v^2}{d(R_{\rm TC})}\,. \tag{12}$$

Therefore, if $\nu_H > 0$ the Higgs becomes lighter relative to $v = 246$ GeV as $N_{\rm TF}$ gets closer to $N_{\rm TF}^c$.

Different model computations have been used to argue whether the ratio $N_{\rm TD}(M_H^0)^2/v^2$ becomes smaller as $N_{\rm TF} \to N_{\rm TF}^c$. These include the technidilaton [19, 21, 22], truncated Schwinger-Dyson equations [14, 37–44], and computations in orientifold-like theories [45, 46]. Perturbative determinations of the conformal window have also shown to lead to a calculable dilaton state parametrically lighter than the other states in the theory [30, 47, 48].

Albeit these reductions from walking dynamics are welcome, Fig. 2 shows that one does not need a large suppression of the TC Higgs (dilaton) mass from walking dynamics when the SM radiative corrections are taken into account.

IV. CONCLUSIONS

We have argued that the observed Higgs mass at the LHC can be interpreted as a TC Higgs with a TeV scale dynamical mass. This is so since the SM top-induced radiative corrections reduce the TC Higgs dynamical mass towards the observed value. We used the phenomenologically motivated assumption that the TC Higgs coupling to the top quark is close to the SM value. In this scenario the next non-Goldstone mesons to be discovered at the LHC have a mass of the order of two to three TeV.

We then investigated the general conditions for the TC spectrum to feature the isosinglet scalar to be identified with the TC Higgs. We used geometric scaling laws, coming from modifying the technicolor matter representation, number of technicolors, techniflavors as well as the number of doublets gauged under the electroweak theory. Finally we also reviewed the potential effects on the TC Higgs properties from walking dynamics and argued that they do not need to be large for achieving a phenomenologically viable TC Higgs physical mass once the SM radiative corrections are taken into account.

Acknowledgments

The work of R.F. is supported by the Marie Curie IIF grant proposal 275012

---

[1] G. Aad *et al.* (ATLAS Collaboration), (2012), arXiv:1207.7214 [hep-ex].
[2] S. Chatrchyan *et al.* (CMS Collaboration), Phys.Lett.B (2012), arXiv:1207.7235 [hep-ex].
[3] M. T. Frandsen and F. Sannino, (2012), arXiv:1203.3988 [hep-ph].
[4] B. Coleppa, K. Kumar, and H. E. Logan, (2012), arXiv:1208.2692 [hep-ph].
[5] E. Eichten, K. Lane, and A. Martin, (2012), arXiv:1210.5462 [hep-ph].
[6] S. Weinberg, Phys.Rev. **D13**, 974 (1976).
[7] S. Dimopoulos and L. Susskind, Nucl.Phys. **B155**, 237 (1979).
[8] E. Eichten and K. D. Lane, Phys.Lett. **B90**, 125 (1980).
[9] F. Sannino, (2008), arXiv:0804.0182 [hep-ph].
[10] F. Sannino, Acta Phys.Polon. **B40**, 3533 (2009), arXiv:0911.0931 [hep-ph].
[11] M. Harada, F. Sannino, and J. Schechter, Phys.Rev. **D54**, 1991 (1996), arXiv:hep-ph/9511335 [hep-ph].
[12] B. Holdom, Phys.Rev. **D24**, 1441 (1981).
[13] T. Appelquist and F. Sannino, Phys.Rev. **D59**, 067702 (1999), arXiv:hep-ph/9806409 [hep-ph].
[14] M. Kurachi and R. Shrock, JHEP **0612**, 034 (2006), arXiv:hep-ph/0605290 [hep-ph].
[15] F. Sannino, Phys.Rev. **D82**, 081701 (2010), arXiv:1006.0207 [hep-lat].
[16] F. Sannino, Phys.Rev.Lett. **105**, 232002 (2010), arXiv:1007.0254 [hep-ph].
[17] S. Di Chiara, C. Pica, and F. Sannino, Phys.Lett. **B700**, 229 (2011), arXiv:1008.1267 [hep-ph].
[18] R. Foadi and F. Sannino, (2012), arXiv:1207.1541 [hep-ph].
[19] K. Yamawaki, M. Bando, and K.-i. Matumoto, Phys.Rev.Lett. **56**, 1335 (1986).
[20] M. Bando, K.-i. Matumoto, and K. Yamawaki, Phys.Lett. **B178**, 308 (1986).
[21] D. D. Dietrich, F. Sannino, and K. Tuominen, Phys.Rev. **D72**, 055001 (2005), arXiv:hep-ph/0505059 [hep-ph].
[22] T. Appelquist and Y. Bai, Phys.Rev. **D82**, 071701 (2010), arXiv:1006.4375 [hep-ph].
[23] J. Beringer *et al.* (Particle Data Group), Phys.Rev. **D86**, 010001 (2012).
[24] F. Sannino, Phys.Rev. **D79**, 096007 (2009), arXiv:0902.3494 [hep-ph].
[25] M. Mojaza, C. Pica, T. A. Ryttov, and F. Sannino, (2012), arXiv:1206.2652 [hep-ph].
[26] D. D. Dietrich and F. Sannino, Phys.Rev. **D75**, 085018 (2007), arXiv:hep-ph/0611341 [hep-ph].
[27] F. Sannino and J. Schechter, Phys.Rev. **D76**, 014014 (2007), arXiv:0704.0602 [hep-ph].
[28] F. Sannino, (2012), arXiv:1205.4246 [hep-ph].
[29] P. de Forcrand, S. Kim, and W. Unger, (2012), arXiv:1208.2148 [hep-lat].
[30] O. Antipin, M. Mojaza, and F. Sannino, (2012), arXiv:1208.0987 [hep-ph].

[31] V. Miransky, Nuovo Cim. **A90**, 149 (1985).

[32] V. Miransky and K. Yamawaki, Mod.Phys.Lett. **A4**, 129 (1989).

[33] V. Miransky and K. Yamawaki, Phys.Rev. **D55**, 5051 (1997), arXiv:hep-th/9611142 [hep-th].

[34] O. Antipin, S. Di Chiara, M. Mojaza, E. Molgaard, and F. Sannino, Phys.Rev. **D86**, 085009 (2012), arXiv:1205.6157 [hep-ph].

[35] P. de Forcrand, M. Pepe, and U. Wiese, Phys.Rev. **D86**, 075006 (2012), arXiv:1204.4913 [hep-lat].

[36] D. Nogradi, JHEP **1205**, 089 (2012), arXiv:1202.4616 [hep-lat].

[37] V. Gusynin and V. Miransky, Sov.Phys.JETP **68**, 232 (1989).

[38] B. Holdom and J. Terning, Phys.Lett. **B187**, 357 (1987).

[39] B. Holdom and J. Terning, Phys.Lett. **B200**, 338 (1988).

[40] M. Harada, M. Kurachi, and K. Yamawaki, Phys.Rev. **D68**, 076001 (2003), arXiv:hep-ph/0305018 [hep-ph].

[41] A. Doff, A. Natale, and P. Rodrigues da Silva, Phys.Rev. **D77**, 075012 (2008), arXiv:0802.1898 [hep-ph].

[42] A. Doff and A. Natale, Phys.Lett. **B677**, 301 (2009), arXiv:0902.2379 [hep-ph].

[43] A. Doff, A. Natale, and P. Rodrigues da Silva, Phys.Rev. **D80**, 055005 (2009), arXiv:0905.2981 [hep-ph].

[44] A. Doff and A. Natale, Phys.Rev. **D81**, 095014 (2010), arXiv:0912.1003 [hep-ph].

[45] F. Sannino and M. Shifman, Phys.Rev. **D69**, 125004 (2004), arXiv:hep-th/0309252 [hep-th].

[46] D. K. Hong, S. D. Hsu, and F. Sannino, Phys.Lett. **B597**, 89 (2004), arXiv:hep-ph/0406200 [hep-ph].

[47] B. Grinstein and P. Uttayarat, JHEP **1107**, 038 (2011), arXiv:1105.2370 [hep-ph].

[48] O. Antipin, M. Mojaza, and F. Sannino, Phys.Lett. **B712**, 119 (2012), arXiv:1107.2932 [hep-ph].